

3. The UK Small Aneurysm Trial Participants. Mortality results for randomised controlled trial of early elective surgery or ultrasonographic surveillance for small abdominal aortic aneurysms. *Lancet* 1998;352:1649-55.
4. Sonesson B, Sandgren T, Lanne T. Abdominal aortic aneurysm wall mechanics and their relation to risk of rupture. *Eur J Vasc Endovasc Surg* 1999;18:487-93.
5. Darling RC, Messina CR, Brewster DC, Ottinger LW. Autopsy study of unoperated abdominal aortic aneurysms. The case for early resection. *Circulation* 1977;56(3 suppl):III161-4.
6. Nicholls SC, Gardner JB, Meissner MH, Johansen HK. Rupture in small abdominal aortic aneurysms. *J Vasc Surg* 1998;28:884-8.
7. Ashton HA, Buxton MJ, Day NE, Kim LG, Marteau TM, Scott RA, et al. The Multicentre Aneurysm Screening Study (MASS) into the effect of abdominal aortic aneurysm screening on mortality in men: a randomised controlled trial. *Lancet* 2002;360:1531-9.
8. Vorp DA, Raghavan ML, Webster MW. Mechanical wall stress in abdominal aortic aneurysm: Influence of diameter and asymmetry. *J Vasc Surg* 1998;27:632.
9. Mower WR, Baraff LJ, Sneyd J. Stress distribution in vascular aneurysms: factors affecting risk of aneurysm rupture. *J Surg Res* 1993;55:155-61.
10. Raghavan ML, Vorp DA, Federle MP, Makaroun MS, Webster MW. Wall stress distribution on three-dimensionally reconstructed models of human abdominal aortic aneurysm. *J Vasc Surg* 2000;31:760-9.
11. Stringfellow MM, Lawrence PF, Stringfellow RG. The influence of aorta-aneurysm geometry upon stress in the aneurysm wall. *J Surg Res* 1987;42:425-33.
12. Fagan MJ. Finite element analysis: theory and practice. Harlow, Essex, England: Longman Scientific & Technical, 1992.
13. Venkatasubramanian AK, Mehta T, Fagan MJ, Mylankal KJ, Kuhan G, Ray B, et al. A comparative study of aortic wall stress using finite element analysis for ruptured and non ruptured abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg* 2004;28:168-76.
14. Fillinger MF, Marra SP, Raghavan ML, Kennedy FE. Prediction of rupture risk in abdominal aortic aneurysm during observation: wall stress versus diameter. *J Vasc Surg* 2003;37:724-32.
15. Raghavan ML, Vorp DA. Towards a biomechanical tool to evaluate rupture potential of abdominal aortic aneurysm: identification of a finite strain constitutive model and evaluation of its applicability. *J Biomech* 2000;33:475-82.
16. Bland M. An introduction to medical statistics. Oxford, UK: Oxford University Press; 2000.
17. Nunnally JC. Psychometric theory. New York, NY: McGraw-Hill; 1978.
18. Di Martino E, Bohra A, Vande Geest JP, Gupta N, Makaroun MS, Vorp DA. Biomechanical properties of ruptured versus electively repaired abdominal aortic aneurysm wall tissue. *J Vasc Surg* 2006;43:570-6.
19. Vande Geest JP, Di Martino ES, Bohra A, Makaroun MS, Vorp DA. A biomechanics based rupture potential index for abdominal aortic aneurysm risk assessment; demonstrative application. *Ann N Y Acad Sci* 2006;1085:11-21.
20. Scoti CM, Shkolnik AD, Muluk SC, Finol EA. Fluid structure interaction in abdominal aortic aneurysms: effect of asymmetry and wall thickness. *Biomed Eng Online* 2005;4:64.
21. Mower WR, Quinones WJ, Gambhir SS. Effect of intraluminal thrombus on aortic aneurysm wall stress. *J Vasc Surg* 1997;26:602-8.
22. Wang DH, Makaroun MS, Webster MW, Vorp DA. Effect of intraluminal thrombus in patient specific models of abdominal aortic aneurysm. *J Vasc Surg* 2002;36:598-604.
23. Schurink GW, van Baalen JM, Visser MJ, van Bockel JH. Thrombus within an aortic aneurysm does not reduce pressure on the aneurysm wall. *J Vasc Surgery* 2000;31:501-6.
24. Vorp DA and Vande Geest JP. Biomechanical determinants of abdominal aortic aneurysm rupture. *Arterioscler Thromb Vasc Biol* 2005;25:1558-66.
25. Vande Geest JP, Sachs MS, Vorp DA. A planar biaxial constitutive relation for the luminal layer of intraluminal thrombus in abdominal aortic aneurysms. *J Biomech* 2006;39:2347-54.
26. Vande Geest JP, Sachs MS, Vorp DA. The effects of aneurysm on the biaxial mechanical behaviour of human abdominal aorta. *J Biomech* 2006;39:1324-34.
27. Vorp DA. Biomechanics of abdominal aortic aneurysm. *J Biomech* 2007;40:1806-12.

Submitted Apr 18, 2007; accepted Sep 4, 2007.

INVITED COMMENTARY

Mark F. Fillinger, MD, Lebanon, NH

Heng et al should be congratulated on a well-done study. In addition to validating earlier work by their laboratory and others using similar methods, they have presented important information about interobserver and intraobserver variability. Although one would like to see even lower interobserver variability, it should be emphasized that the authors have explored a "worst-case scenario" in which three of four observers had little training or experience in the method.

Two laboratories now have large clinical series including elective and ruptured abdominal aortic aneurysms (AAAs). Both laboratories have validated that peak wall stress is superior to maximum diameter for estimating rupture risk by a second patient cohort.¹⁻³ There are, of course, methodologic issues that can be improved. Thresholds for determining low and high risk of rupture differ in the laboratories performing these studies, most likely owing to differing methods of creating the mesh. This demonstrates the importance of consistency in the methods used to create the finite element mesh, and the importance of clinical control series to validate thresholds indicating elevated risk for rupture. Other ongoing areas of investigation include the relative impor-

tance of wall strength, calcification, thrombus, and fluid-structure interactions.

Large multi-institutional collaborative efforts are already under way to perform external validation cohorts for the method. A significant amount of work still remains to make these methods widely available to clinicians, but Heng et al should be congratulated on an important step along that pathway.

REFERENCES

1. Fillinger MF, Raghavan ML, Marra SP, Cronenwett JL, Kennedy FE. In vivo analysis of mechanical wall stress and abdominal aortic aneurysm rupture risk. *J Vasc Surg* 2002;36:589-97.
2. Fillinger MF, Marra SP, Raghavan ML, Kennedy FE. Prediction of rupture risk in abdominal aortic aneurysm during observation: wall stress versus diameter. *J Vasc Surg* 2003;37:724-32.
3. Venkatasubramanian AK, Fagan MJ, Mehta T, Mylankal KJ, Ray B, Kuhan G, et al. A comparative study of aortic wall stress using finite element analysis for ruptured and non-ruptured abdominal aortic aneurysms. *Eur J Vasc Endovasc Surg* 2004;28:168-76.